NAME

DATE

PERIOD

Unit 8, Lesson 3 Rational and Irrational Numbers

Let's learn about irrational numbers.


3.1 Algebra Talk: Positive Solutions

Find a positive solution to each equation:

1. $x^2 = 36$

- 2. $x^2 = \frac{9}{4}$
- 3. $x^2 = \frac{1}{4}$
- 4. $x^2 = \frac{49}{25}$

3.2 Three Squares



1. Draw 3 squares of different sizes with vertices aligned to the vertices of the grid.

NAME	DATE	PERIOD
2. For each square:		
a. Label the area.		

- b. Label the side length.
- c. Write an equation that shows the relationship between the side length and the area.

3.3 Looking for a Solution

Are any of these numbers a solution to the equation $x^2 = 2$? Explain your reasoning.

- 1
- •
- $\frac{1}{2}$ $\frac{3}{2}$
- $\frac{7}{5}$

3.4 Looking for $\sqrt{2}$

A rational number is a fraction or its opposite (or any number equivalent to a fraction or its opposite).

1. Find some more rational numbers that are close to $\sqrt{2}$.

2. Can you find a rational number that is exactly $\sqrt{2}$?

NAME

DATE

PERIOD

♣ Are you ready for more?

If you have an older calculator evaluate the expression

 $\left(\frac{577}{408}\right)^2$

and it will tell you that the answer is 2, which might lead you to think that $\sqrt{2} = \frac{577}{408}$.

- 1. Explain why you might be suspicious of the calculator's result.
- 2. Find an explanation for why $408^2 \cdot 2$ could not possibly equal 577^2 . How does this show that $\left(\frac{577}{408}\right)^2$ could not equal 2?
- 3. Repeat these questions for

$$\left(\frac{1414213562375}{100000000000}\right)^2 \neq 2,$$

an equation that even many modern calculators and computers will get wrong.

NAME


PERIOD

DATE

Lesson 3 Summary

In an earlier activity, we learned that square root notation is used to write the length of a side of a square given its area. For example, a square whose area is 2 square units has a side length of $\sqrt{2}$ units, which means that

$$\sqrt{2} \cdot \sqrt{2} = 2.$$



A square whose area is 25 square units has a side length of $\sqrt{25}$ units, which means that

$$\sqrt{25} \cdot \sqrt{25} = 25$$

Since $5 \cdot 5 = 25$, we know that

$$\sqrt{25} = 5.$$

 $\sqrt{25}$ is an example of a rational number. A **rational number** is a fraction or its opposite. Remember that a fraction $\frac{a}{b}$ is a point on the number line found by dividing the segment from 0 to 1 into *b* equal intervals and going *a* of those intervals to the right of 0. We can always write a fraction in the form $\frac{a}{b}$ where *a* and *b* are whole numbers (and *b* is not 0),

but there are other ways to write them. For example, we can write $\sqrt{25} = \frac{5}{1}$. You first learned about fractions in earlier grades, and at that time, you probably didn't know about negative numbers. Rational numbers are fractions, but they can be positive or negative. So, -5 is also a rational number. Because fractions and ratios are closely related ideas, fractions and their opposites are called RATIOnal numbers.


Here are some examples of rational numbers:

$$\frac{7}{4}, 0, \frac{6}{3}, 0.2, -\frac{1}{3}, -5, \sqrt{9}, -\frac{\sqrt{16}}{\sqrt{100}}$$

Can you see why they are each examples of "a fraction or its opposite?" An **irrational number** is a number that is not rational. That is, it is a number that is not a fraction or its opposite. $\sqrt{2}$ is an example of an irrational number. It has a location on the number line, and its location can be approximated by rational numbers (it's a tiny bit to

NAME	DATE	PERIOD

the right of $\frac{7}{5}$), but $\sqrt{2}$ can not be found on a number line by dividing the segment from 0 to 1 into *b* equal parts and going *a* of those parts away from 0 (if *a* and *b* are whole numbers).



 $\frac{17}{12}$ is also close to $\sqrt{2}$, because $\left(\frac{17}{12}\right)^2 = \frac{289}{144}$. $\frac{289}{144}$ is very close to 2, since $\frac{288}{144} = 2$. But we could keep looking forever for solutions to $x^2 = 2$ that are rational numbers, and we would not find any. $\sqrt{2}$ is not a rational number! It is irrational.

In your future studies, you may have opportunities to understand or write a proof that $\sqrt{2}$ is irrational, but for now, we just take it as a fact that $\sqrt{2}$ is irrational. Similarly, the square root of any whole number is either a whole number ($\sqrt{36} = 6$, $\sqrt{64} = 8$, etc.) or irrational ($\sqrt{17}$, $\sqrt{65}$, etc.). Here are some other examples of irrational numbers:

$$\sqrt{10}, -\sqrt{3}, \frac{\sqrt{5}}{2}, \pi$$

Glossary Terms

irrational number

rational number